

# Bio-potential application of algae in reducing the toxicity of the fly ash from coal processing: a new trend of reclamation

## Abstract

Fresh water algal growth inhibition test with unicellular alga *Ankistrodesmus falcatus*, *Scenedesmus obliquus*, *Selenastrum capricornutum*, and *Microcoleus vaginatus* is of great interest in biodiversity science. Our investigation has been made in laboratory continuous culture using sterile sewage as the medium for the start-up growth of these species. And since, these are at risk level when meet the coal industry waste, a suitable bioassay application based on an oxidizing medium of arsenic, cadmium, mercury, and selenium was implemented for algal incubation. First, the algistatic-algital levels were reported. Second, the observed growth inhibition with the value of IC 50 (the concentration which inhibits algae growth by 50%) was investigated to obtain the growth responses and to define the range of the studied elemental variations. Noticeably, the median EC50 of the scrubber ash slurry extract (SASE) was 3-15%. Further, the results showed that the toxicity of the heavy metals depends on their chemical speciation and can be related to their free ion activities and their concentrations.

**Keywords:** toxicity, fly ash, *Microcoleus vaginatus*, oxidizing medium, EC50, ecosystem

Volume 2 Issue 2 - 2018

Nuha Al Masri

Department of Chemistry, Syrian Private University, Syria

**Correspondence:** Noha Almasri, Department of Chemistry, Faculty of Medicine, Syrian Private University, Damascus, Syria, Tel +963 9334 46993, Email [almasrinuhasy@gmail.com](mailto:almasrinuhasy@gmail.com)

**Received:** March 06, 2018 | **Published:** April 20, 2018

## Introduction

Increasing energy demands and the national program emphasizing the use of coal have stimulated mining and on-site combustion of vast deposits in several countries. These activities have posed new environmental considerations in many regions previously undisturbed or even pristine in character. Bituminous and sub-bituminous coals contain a broad spectrum of trace-elements, some of which have toxic properties (e.g. arsenic, cadmium, mercury, and selenium) and have been shown to be considerably enriched in coal relative to average crustal abundance on Earth.<sup>1</sup> Additional enrichment of the more volatile toxic elements occurs during the combustion process; subsequent condensation takes place on the surface of submicron fly ash particles which escape emission control devices and become biologically available.<sup>2</sup>

Present emission control technology, which can exceed 98% removal of fly ash particulates, includes electrostatic precipitators and venturi wet scrubbers. Both types of installations are found in the newer, large western coal-fired power plants.<sup>3</sup> Both electrostatic precipitators and wet venturi scrubber systems used in coal-fired generating plants produce an ash that is also a potential pollutant. Mahlaba et al.<sup>4</sup> indicated that molybdate, borate, fluoride, selenate, and arsenate were soluble contaminants from coal ash disposed in alkaline environments.

The potential deposition of trace elements from coal-fired generating plants has received increasing attention in recent years due to their adverse effects on the environment.<sup>5</sup> To minimize the effects of these toxic trace elements, their deposition rates should not reach concentrations which are incompatible with naturally functioning ecosystems. The compatible level for each trace element depends on the interaction of physicochemical and biological components of the ecosystem which directly or indirectly determine the chemical state

of the element, its concentration, its availability,<sup>6</sup> and the potential for bioconcentration in the food web.

The use of algal toxicity tests to monitor the biological impact of pollutants is important because algae are primary producers in aquatic systems. Such tests provide one approach to determining compatible levels of trace elements, although care must be taken to create a test situation which represents as closely as possible the natural situation being examined. Because pollutants from coal-fired generating plants typically enter aquatic ecosystems in small quantities over long periods,<sup>6</sup> a chronic exposure test is indicated when investigating their potential toxicity.

The objectives of this study were to develop an algal toxicity test system, based on the EPA Algal Assay Procedure Bottle Test, for determining the median effective concentration (EC50) values of heavy metals as potential pollutants from coal-fired power plants.<sup>7</sup> The EC50 values which have been developed for As(V), Cd(II), Hg(II), and Se(VI), in solution, and for extracts of scrubber ash slurry from a western coal-fired generating plant are based on dose-response relationships for algae indigenous to southeastern part of Syria and *Selenastrum capricornutum*, an EPA test algae.

## Materials and methods

The Algal Assay Procedure Bottle Test (AAPBT) was adapted to evaluate the effect of As(V), Cd(II), Hg(II), and Se(VI), in solution, and of scrubber ash slurry extract from the Colstrip coal-fired power generating plant on selected algal species. The algae used in these investigations were the freshwater chlorophytes *Ankistrodesmus falcatus* (CoMa) Ralfs, *Scenedesmus obliquus* (Turp.) Kutz., and *Selenastrum capricornutum* Printz, and the freshwater cyanophyte *Microcoleus vaginatus* (Vauch.) Gom. These species subsequently were referred to by their generic names.

*Ankistrodesmus*, *Scenedesmus*, and *Microcoleus* were isolated from phytoplankton collections from stock ponds in Sweda, Syria. The unialgal, bacteria-free cultures of *Ankistrodesmus* and *Scenedesmus* were obtained by repeated streaking of isolated colonies on Petri plates containing 1.5% Bacto-Agar (Difco) in Algal Assay Medium (AAM).<sup>8</sup> Stock cultures of *Ankistrodesmus* and *Scenedesmus* were maintained in AAM. *Microcoleus* was isolated using similar streaking techniques but substituting Gorham's nutrient medium<sup>9</sup> for AAM. Stock cultures of *Microcoleus* were grown in Gorham's medium because *Microcoleus* could not be maintained in AAM. The unialgal *Microcoleus* was not bacteria-free. An axenic culture of *Selenastrum* was obtained and maintained in AAM. The axenic cultures were checked for bacteria with each transfer using Difco Tryptone Glucose Extract Agar (TGEA).

Laboratory toxicity testing was followed procedures outlined by the Syrian Public Health Association. Polycarbonate flasks and all glassware in contact with trace element solutions or scrubber ash slurry extract were rinsed with 10% HNO<sub>3</sub> solutions. All other glassware was rinsed with recommended 10% HCl solutions.

Polycarbonate flasks, rather than glass, were used for toxicity test containers in the trace element toxicity studies. Some investigators have used polycarbonate flasks.<sup>10</sup> Hai et al.<sup>11</sup> reported that mercury adsorption on the walls of silicone glass was a serious problem. Lee et al.<sup>12</sup> found adsorption of cadmium on the walls of Pyrex glass also was a cause for concern. Investigations with <sup>76</sup>As(III), <sup>115m</sup>Cd(II), and <sup>75</sup>Se(IV) in our laboratory indicated that less than 5% of the elements were adsorbed on the walls of polycarbonate flasks at 0.01, 0.001 and 0.01mg/L, respectively.

Similar test conditions were imposed for all toxicity tests. The culture media used for toxicity tests were the same as those used for stock cultures. The chelator (EDTA) was omitted in toxicity tests, although small quantities necessarily were introduced via the algal inocula. The micronutrients also were omitted from AAM. Chemical precipitates remaining in the culture media after autoclaving were dissolved by the addition of 1N HCl. The test elements were introduced as analytical reagent grade salts of Na<sub>2</sub>HAsO<sub>4</sub>·H<sub>2</sub>O, CdSO<sub>4</sub>, HgSO<sub>4</sub>, and Na<sub>2</sub>SeO<sub>4</sub>. Stock solutions containing 1g/L element were prepared using deionized water. The stock solutions were acidified with HCl to pH 1.5. The appropriate range of test concentrations was obtained by diluting the stock solutions aseptically with sterile culture media. Element levels of 0 (control), 0.01, 0.1, 1, 10 and 100 mg/L were used in the initial toxicity screenings. Treatment levels of subsequent experiments were dependent on initial results of toxicity screening. The pH of each test culture solution was adjusted to 7.0 ± 0.3 with NaOH or HCl solution prior to inoculation with the test alga.

The toxicity tests with scrubber ash slurry were conducted using an extract of the scrubber ash slurry obtained from the settling pond for Units 1 and 2 of the coal-fired generating plant. The extract was prepared by filtering the slurry (10% solids) through 0.45µm Millipore filters, mixing the filtrate with the appropriate algal medium, and filter-sterilizing through 0.22µm Millipore filters. The toxicity tests were conducted using percentages of algal medium/slurry extract stock solutions in the appropriate algal medium. The pH of the test solution varied with the concentration of the slurry extract. The scrubber ash slurry extract (SASE), without nutrients, was analyzed for pH, specific conductance, alkalinity, total phosphate, ammonia nitrogen, nitrate-nitrite nitrogen, total organic carbon, silica, and sulfate. An aliquot also was analyzed for selected trace elements using

inductively coupled plasma-atomic emission spectroscopy.

All treatment levels of the trace elements and SASE were tested in triplicate for each experimental run. The experimental runs were repeated for each combination until the dose-response relationship was well-defined. The number of runs ranged from two to five for each experimental combination. Test flasks were inoculated from 10- to 14-day axenic cultures of *Ankistrodesmus*, *Scenedesmus*, or *Selenastrum* to give a final cell concentration of 1 x 10<sup>4</sup> cells/mL ± 10%. The *Microcoleus* inoculum, of the same age, was 1mL of a stock culture reading 30% transmission at 450nm on a Hach DR-EL2 spectrophotometer. All test cultures were incubated in 500-mL polycarbonate flasks containing 100ml of test culture solution. The cultures were incubated at 24±2°C under 400ft-c±10% mixed illumination ("cool white" fluorescent and incandescent), measured adjacent to the flask at the liquid level, on a 16:8 h light: dark regime.

After two weeks the algae were harvested and analyzed for chlorophyll a (Trichromatic Method) as outlined by the Syrian Public Health Association. The extracted chlorophyll was read on a modified Gilford spectrophotometer with a 1-cm light path. Percent of control (response) values were calculated using chlorophyll a values. The chlorophyll a detection limit of the algal assay method was approximately the same as the chlorophyll a inoculation level. Therefore, either an algistatic or algicidal response could be indicated by a chlorophyll a value of zero. A t-test was used to determine if the response was significantly different from 100% of the control. Concentrations of the toxic solutions which produced 50% of the control responses, i.e. the median effective concentrations (EC50 values), were found by fitting a line to the dose-response relationship by least squares using the log<sub>10</sub> of the dose as the independent variable.

## Results

The growth responses (% of control values) of *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus* on exposure to As(V), Cd(II), Hg(II), Se(VI), and SASE are presented in Tables 1-5, respectively. *Selenastrum* is increasingly inhibited with an increasing As(V) concentration between 10 and 75mg/L. A similar response was found for *Ankistrodesmus* and *Scenedesmus*, from 0.01 to 5 and 0.01 to 50mg/L, respectively. The first treatment level to cause a statistically significant inhibitory response to *Selenastrum* was 25 mg/L. The first significant growth inhibition of *Ankistrodesmus* and *Scenedesmus* occurred at 0.1 and 0.01 mg/L AS(V), respectively. *Microcoleus* was not significantly inhibited at any treatment level. In fact, a stimulatory response was noted at 75 mg/L. Algistatic-algicidal responses were noted for *Ankistrodesmus* and *Scenedesmus* at 5 and 50mg/L As(V), respectively.

Cadmium (II) was extremely inhibitory, causing significant inhibition of *Ankistrodesmus*, *Scenedesmus*,

and *Microcoleus* at 0.01mg/L and *Selenastrum* at 0.05 mg/L (Table 2). All the algae responded algistatically-algicidally at or below 0.3mg/L. A significant stimulatory response was noted for *Ankistrodesmus* with 0.001mg/L.

The sensitivity of the algal species to Hg (II) was variable (Table 3). The first significant inhibition of *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus* occurred at 0.05, 0.1, 0.01 and 0.4 mg/L Hg (II), respectively. Algistatic-algicidal responses for *Ankistrodesmus*, *Scenedesmus*, and *Selenastrum* were first noted at 0.4mg/L. *Microcoleus* approached an algistatic response at 1.0mg/L.

The sensitivity of the algal species to Se (VI) varied (Table 4). The first significant inhibition of *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus* occurred at 0.01, 0.1, 0.3 and 10 mg/L Se (VI), respectively. *Microcoleus* was stimulated by treatment levels of 1.0 mg/L and below. Algistatic-algicidal responses by *Ankistrodesmus*, *Scenedesmus*, and *Selenastrum* appeared at 10, 4 and 1.7 mg/L, respectively. *Microcoleus* approached an algistatic response at 50mg/L.

The first significant inhibition of *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus* occurred at 1, 5, 10 and 5% SASE, respectively (Table 5). Algistatic-aigicidal responses for *Ankistrodesmus*, *Scenedesmus*, and *Selenastrum* were indicated at 50, 100 and 75% SASE. *Microcoleus* approached an algistatic response

at 100% SASE.

The chemical analysis of the SASE is presented in Table 6. The extract was high in sulfate, nitrogen (nitrate-nitrite), total phosphate, silica, calcium, magnesium, manganese, specific conductance and alkalinity. Several potentially toxic trace elements were found to be present at low levels.

The regression models used for predicting EC50 values of As (V), Cd (II), Hg (II), Se (VI), and SASE are presented in Table 7. No regression model is presented for the As (V)-*Microcoleus* combination because *Microcoleus* was not significantly inhibited at the highest treatment level. All of the final regression models except Hg (II)-*Scenedesmus* and Se(VI)-*Scenedesmus* have coefficient of determination (R<sup>2</sup>) values above 0.60 (Table 7).

**Table 1** Growth responses of *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus* to As(V)

As(V), mg/L	<i>Ankistrodesmus</i>		<i>Scenedesmus</i>		<i>Selenastrum</i>		<i>Microcoleus</i>	
	n <sup>a</sup>	y±se <sup>b</sup>	n	y±se	n	y±se	n	y±se
0.01	3	61±18	5	58±20*	6	78±26	6	106±23
0.1	3	72±4*	6	48±15*	6	87±30	6	97±13
1	6	52±22*	6	26±10*	6	76±24	6	99±17
5	3	0	-	-	-	-	-	-
10	6	0	5	4±5*	5	104±26	6	106±13
25	3	0	3	1±2*	3	50±18*	3	90±17
50	6	0	3	0	3	9±2*	3	78±26
75	6	0	3	0	3	4±10*	3	115±4*
100	6	0	3	0	6	14±15*	6	97±17

<sup>a</sup>Number of observations.

<sup>b</sup>Standard error of the mean.

\*Value significantly different from 100% (p<0.05).

**Table 2** Growth responses of *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus* to Cd(II)

Cd(II), mg/L	<i>Ankistrodesmus</i>		<i>Scenedesmus</i>		<i>Selenastrum</i>		<i>Microcoleus</i>	
	n <sup>a</sup>	y±se <sup>b</sup>	n	y±se	n	y±se	n	y±se
0.001	3	156±13*	2	118±11	2	89±73	3	118±23
0.005	3	99±34	3	78±18	3	106±58	3	89±21
0.01	7	69±29*	8	8±12*	6	61±28	6	69±12*
0.05	6	8±5*	6	0	6	17±7*	6	3±3*
0.1	9	1±1*	9	0	8	7±5*	6	1±1*
0.2	3	0	3	0	3	2±1*	-	-
0.3	6	0	6	0	6	0	3	0

<sup>a</sup>Number of observations.

<sup>b</sup>Standard error of the mean.

\*Value significantly different from 100% (p<0.05).

The EC50 values for the test algae exposed to the trace elements and SASE are presented in Table 8. The EC50 values ranged from 0,048-30.761 mg/L (0.00064-0.41058 M) As (V), 0.005-0.019mg/L (0.00004-0.00017 M) Cd(II), 0.033-0.253mg/L (0.00016--0.00126

M) Hg (II), 0.033-8.511mg/L (0.00042-0.10779 M) Se(VI), and 3.048-15.417% SASE. The test algae with the lowest EC50 values were *Scenedesmus* for As (V) and Cd (II), *Selenastrum* for Hg(II), *Ankistrodesmus* for Se(VI), and *Microcoleus* for SASE.

**Table 3** Growth responses of *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus* to Hg(II)

Hg (II), mg/L	<i>Ankistrodesmus</i>		<i>Scenedesmus</i>		<i>Selenastrum</i>		<i>Microcoleus</i>	
	n <sup>a</sup>	y±se <sup>b</sup>	n	y±se	n	y±se	n	y±se
0.001	3	91±28	6	150±78	3	99±12	-	-
0.01	5	92±14	9	86±19	6	88±11*	6	99±10
0.05	3	70±11*	5	86±39	3	52±21	5	79±40
0.1	8	48±32*	8	55±46*	6	6±10*	6	102±12
0.4	3	0	6	0	3	0	5	42±22*
0.7	3	0	6	0	3	0	5	12±16*
1	9	0	9	0	9	0	5	1±2*

<sup>a</sup>Number of observations.

<sup>b</sup>Standard error of the mean.

\*Value significantly different from 100% (p<0.05).

**Table 4** Growth responses of *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus* to Se(IV)

Se(IV), mg/L	<i>Ankistrodesmus</i>		<i>Scenedesmus</i>		<i>Selenastrum</i>		<i>Microcoleus</i>	
	n <sup>a</sup>	y±se <sup>b</sup>	n	y±se	n	y±se	n	y±se
0.01	7	70±31*	3	236±129	3	89±8	3	118±1*
0.05	3	46±1*	6	152±70	-	-	3	105±12
0.1	9	30±14*	15	48±32*	8	84±19	3	116±9
0.2	3	19±2*	3	58±15*	3	81±19	-	-
0.3	3	18±3*	3	21±5*	3	48±17*	-	-
0.4	3	10±4*	8	19±10*	6	20±14*	3	128±6*
0.5	3	13±3*	-	-	3	32±11*	-	-
0.6	3	10±2*	-	-	2	33±5*	-	-
0.7	3	9±2*	7	18±8*	6	16±8*	3	117±7*
0.8	2	12±2*	-	-	3	15±1*	-	-
0.9	3	11±3*	-	-	3	18±5*	-	-
1	8	10±10*	12	12±8*	9	5±6*	6	110±3*
1.4	3	12±0*	2	11±4*	3	2±3*	-	-
1.7	3	7±1*	3	8±3*	3	0	-	-
2	3	8±3*	2	8±8*	3	0	-	-
4	-	-	6	0	-	-	-	-
5	-	-	-	-	-	-	3	73±11
10	3	0	-	-	3	0	3	50±6*
20	-	-	-	-	-	-	2	16±4*
30	-	-	-	-	-	-	3	6±2*
40	-	-	-	-	-	-	3	4±3*
50	-	-	-	-	-	-	3	5±2*

<sup>a</sup>Number of observations.

<sup>b</sup>Standard error of the mean.

\*Value significantly different from 100% (p<0.05).

**Table 5** Growth responses of *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus* to scrubber ash slurry extract (SASE).

SASE	<i>Ankistrodesmus</i>		<i>Scenedesmus</i>		<i>Selenastrum</i>		<i>Microcoleus</i>	
	n <sup>a</sup>	y±se <sup>b</sup>	n	y±se	n	y±se	n	y±se
0.01	6	98±32	5	103±26	4	98±42	5	106±15
0.1	6	72±42	5	72±26	6	88±31	5	103±6
1	6	72±18*	5	115±61	4	135±66	6	98±15
5	6	57±17*	5	65±9*	6	98±42	5	19±4*
10	6	61±15*	5	62±17*	6	61±20*	6	8±3*
25	6	4±3*	6	5±3*	6	40±10*	6	7±1*
50	5	0	6	1±1*	6	2±3*	6	4±4*
75	6	0	6	2±2*	6	0	6	1±2*
100	6	0	6	0	6	0	6	1±1*

<sup>a</sup>Number of observations.

<sup>b</sup>Standard error of the mean.

\*Value significantly different from 100% (p<0.05).

**Table 6** Chemical analysis of scrubber ash slurry extract (SASE)

Parameter	Concentration <sup>a</sup>	Parameter	Concentration <sup>a</sup>
pH	7.3	Ba	0.04
Specific conductance	9750	Be	0
Alkalinity <sup>b</sup>	248	Ca	360
SO <sub>4</sub> <sup>2-</sup>	10.2	Cd	0.04
C-total organic	1	Co	0.32
NH <sub>3</sub> -N	0.4	Cu	0
NO <sub>2</sub> <sup>-</sup> +NO <sub>3</sub> <sup>-</sup> -N	21.4	Fe	0.07
PO <sub>4</sub> <sup>3-</sup> -total	1.6	Mg	>100
Si	91	Mn	47
Al	0.56	Ni	0.3
As	0.25	Zn	0.12

<sup>a</sup>µmhos cm<sup>-1</sup> for specific conductance and mg/L for all other parameters except pH.

<sup>b</sup>pH 4.5 as CaCO<sub>3</sub>.

**Table 7** Regression equations of algal responses to selected trace element or SASE treatment levels

Test substance	Alga	Regression equation	R <sup>2</sup>
As	<i>Ankistrodesmus</i>	Y= -26.963 (log <sub>10</sub> X)+34.053	0.64
	<i>Scenedesmus</i>	Y=-18.700 (log <sub>10</sub> X)+25.251	0.7
	<i>Selenastrum</i>	Y= -96.029 (log <sub>10</sub> X)+192,868	0.79
Cd	<i>Ankistrodesmus</i>	Y=-78.401 (log <sub>10</sub> X)-84.306	0.89
	<i>Scenedesmus</i>	Y=-115.248 (log <sub>10</sub> X)-215.247	0.82
	<i>Selenastrum</i>	Y=-65.792 (log <sub>10</sub> X)-63.788	0.64
	<i>Microcoleus</i>	Y= -65.747 (log <sub>10</sub> X)-70.207	0.91
Hg	<i>Ankistrodesmus</i>	Y= -54.274 (log <sub>10</sub> X)- 10.201	0.63
	<i>Scenedesmus</i>	Y=-50.592 (log <sub>10</sub> X)-4.257	0.46
	<i>Selenastrum</i>	Y=-78.388 (log <sub>10</sub> X)-66.496	0.86
	<i>Microcoleus</i>	Y=-72.355 (log <sub>10</sub> X)+6.822	0.67

Table Continued..

Test substance	Alga	Regression equation	R <sup>2</sup>
Se	<i>Ankistrodesmus</i>	Y= -30.462 (log <sub>10</sub> X)+4.754	0.69
	<i>Scenedesmus</i>	Y= -73.596 (log <sub>10</sub> X)+10.811	0.53
	<i>Selenastrum</i>	Y= -79.893 (log <sub>10</sub> X)+5.465	0.8
	<i>Microcoleus</i>	Y= -66.669 (log <sub>10</sub> X)+112.036	0.96
SASE	<i>Ankistrodesmus</i>	Y= -92.339 (log <sub>10</sub> X)+147.314	0.79
	<i>Scenedesmus</i>	Y=-70.753 (log <sub>10</sub> X)+116.941	0.71
	<i>Selenastrum</i>	Y= -78.803 (log <sub>10</sub> X)+143.654	0.67
	<i>Microcoleus</i>	Y=-93.000 (log <sub>10</sub> X)+94.993	0.93

<sup>a</sup> x is the concentration of the test substance.

**Table 8** EC50 values<sup>a</sup> of the trace elements and SASE for the test algae

Test substance	Alga	Log <sub>10</sub> EC50	EC50 <sup>b</sup>
As	<i>Ankistrodesmus</i>	- 0.59±0.17 <sup>c</sup>	0.256
	<i>Scenedesmus</i>	- 1.32±0.17	0.048
	<i>Selenastrum</i>	1.49±0.20	30.8
Cd	<i>Ankistrodesmus</i>	-1.71±0.12	0.019
	<i>Scenedesmus</i>	-2.30±0.32	0.005
	<i>Selenastrum</i>	- 1.73±0.25	0.019
	<i>Microcoleus</i>	- 1.83±0.13	0.015
Hg	<i>Ankistrodesmus</i>	1.11±0.25	0.078
	<i>Scenedesmus</i>	- 1.07±0.24	0.085
	<i>Selenastrum</i>	- 1.48±0.17	0.033
	<i>Microcoleus</i>	-0.60±0.11	0.253
Se	<i>Ankistrodesmus</i>	- 1.49±0.10	0.033
	<i>Scenedesmus</i>	-0.53±0.09	0.294
	<i>Selenastrum</i>	-0.56±0.03	0.277
SASE	<i>Microcoleus</i>	0.93±0.06	8.51
	<i>Ankistrodesmus</i>	1.05±0.18	11.3
	<i>Scenedesmus</i>	0.95±0.15	8.83
SASE	<i>Selenastrum</i>	1.19±0.16	15.4
	<i>Microcoleus</i>	0.48±0.05	3.05

<sup>a</sup>mg/L trace elements and % SASE.

<sup>b</sup>EC50 re-transformed from log scale.

<sup>c</sup>Standard error of estimate.

## Discussion

Toxicity tests provide an important method of assessing the biological effects of elements in solution, and the adapted Algal Assay Procedure Bottle Test-Algal Assay Medium (AAPBT-AAM) provides a good culture medium for evaluating the effect of dissolved substances on algae. The medium is buffered to maintain pH near neutrality has a total dissolved solids concentration of less than 70mg/L and when formulated without EDTA, is completely inorganic. The importance of using a test medium similar in ionic strength to the lower ionic

strength of natural fresh waters is indicated in the literature.<sup>13</sup>

Chelating agents were omitted from our culture solutions because they have been shown to counteract inhibition of growth in toxicity studies.<sup>14</sup> Awasthi & Das<sup>15</sup> omitted organic chelators to minimize chemical complications when investigating heavy metal toxicity to algae. Additionally, Tawarada et al.<sup>16</sup> reported EDTA displays a strong tendency to complex with cadmium. The adjustment of the algal assay solution to pH 7.0±0.3 with NaOH or HCl reduced the possibility of pH-trace element interactions affecting the algal responses. Laboratory investigations indicated that small additions of NaOH and HCl did not significantly affect growth of the algal species. This methodology was used by Horvatić et al.<sup>17</sup> Vigneault & Campbell<sup>18</sup> reported that cadmium accumulation was pH dependent.

The use of HCl to dissolve particulates formed during autoclaving of the medium reduced the possibility of occlusion, adsorption, or precipitation of the trace element being studied.<sup>19</sup> Pyta & Rogula-Kozłowska<sup>20</sup> found that mercury accumulated in the particulate fraction of the culture medium. Martins et al.<sup>21</sup> speculated that cadmium was occluded when calcium carbonate precipitated in his test solutions. Micronutrients were omitted in order to reduce the possibility of synergistic or antagonistic effects. Hart & Bertram<sup>22</sup> reported that the concentration of manganese in the medium regulated the amount of cadmium accumulated in *Chlorella pyrenoidosa*. Virtually no cadmium accumulation occurred in cells grown in medium containing 0.2mg/L manganese. Wong & Beaver<sup>23</sup> also omitted heavy metal micronutrients from the medium when investigating heavy metal toxicity with algal bioassays. Use of only a small quantity of sulfur in selenium assays is important because sulfur and selenium compounds have been shown to display antagonistic effects.<sup>24</sup>

To provide a viable extrapolation from the laboratory to the natural environment, we selected the dominant oxidation states of the trace elements occurring in unpolluted, natural freshwaters which are aerobic and slightly alkaline. Axenic unialgal cultures were used, when possible, for these static algal toxicity tests. Bacterial conversion of mercury ion to the organic and/or elemental form and subsequent volatilization from non-acidified media was reported by Finley & McLaughlin.<sup>25</sup> Microorganisms vary from species to species in their sensitivity to toxic substances, exhibit different dose-response relationships, and display different bioconcentration abilities. For this reason, considerable variance between experiments should be anticipated with a mixed inoculum because the relative species composition of the inocula would not be consistent. Unknown

interactions could occur, making interexperimental comparisons of dose-response relationships difficult.

The use of an inoculum of uniformly low density in these investigations reduced the possibility of an algal overload. Hart & Bertram<sup>22</sup> indicated that cell concentration affected the toxicity of Hg (II) to *Chlorella pyrenoidosa*. Pearson<sup>26</sup> reported that the toxicity of mercurial compounds decreased with increasing cell concentrations. Variations in the growth phase and metabolic activity of the inoculum were reduced by inoculating the test solutions from 10- to 14-day stock cultures. Prior determination of growth curves for each alga under the test conditions established that cultures of this age were healthy and had not entered the stationary phase, although the population was increasing at a decreasing rate. The AAPBT recommends the use of 10- to 14-day cultures of *Selenastrum capricornutum*. A 2 week period for the algal toxicity studies was chosen for these reasons.

A standard temperature of 24±2°C was used because temperature-trace metal interactions have been shown to affect algal responses.<sup>27</sup> The static toxicity tests employed in this investigation may have tended to underestimate the toxicity of low-level treatments because the low-level concentrations of the potential metal pollutants could have been further reduced during the tests through bioaccumulation and possibly volatilization in the case of Hg (II).<sup>28</sup> If the tests had been conducted in a continuous-flow algal system, then the first significant inhibition levels of As (V), Cd (II), Hg (II), Se (VI), and SASE might have been reduced. The EC50 values, however, are much less affected by this depletion.

Little information is available concerning the toxicity of arsenic to freshwater algae. The algistatic-algicidal value reported for *Chlorella vulgaris* of 0.06mg/L As(V) by Jiang et al.<sup>29</sup> is well below the algistatic-algicidal value of 5.0mg/L As(V) for *Ankistrodesmus*, the alga with the lowest algistatic-algicidal value. Mekhalfi et al.<sup>30</sup> reported no detrimental effects to *Asterionella formosa* when this freshwater diatom was exposed to 0.16mg/L As(V). Additionally, the EC50 values (0.048 to > 100mg/L) and the algistatic-algicidal values (5 to > 100mg/L) displayed a large degree of variation from alga to alga. The blue-green alga *Microcoleus* was much more tolerant of As(V) than the green algae investigated. This could be due in part to differences in the assay media, the fact that the *Microcoleus* was not axenic, or possibly because *Microcoleus* is a prokaryotic organism while the other organisms tested were eukaryotes.

This investigation confirms reports that cadmium is extremely toxic to freshwater algae. Cadmium concentrations of 0.13 and 0.08mg/L for *Selenastrum capricornutum*, 0.14mg/L for *Chlorella vulgaris*, and >0.01mg/L for *Asterionella formosa* were reported as algistatic-algicidal,<sup>31</sup> respectively. In this investigation, algistatic-algicidal values between 0.1-0.2, 0.01-0.05, 0.2-0.3 and 0.1-0.3mg/L Cd(II) were indicated for *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus*, respectively.

Hart & Bertram<sup>22</sup> reported that a Cd(II) concentration of 0.25mg/L as Cd(CH<sub>3</sub>CO<sub>2</sub>)<sub>2</sub>·2H<sub>2</sub>O inhibited the growth rate of *Chlorella pyrenoidosa* cultures in the logarithmic growth phase. Cheng et al.<sup>31</sup> found *Chlorella vulgaris* to be more sensitive and reported an abrupt inhibition of growth by concentrations of cadmium above 0.05mg/L. Liu et al.<sup>32</sup> observed that Cd(II) concentrations as low as 0.0061 mg/L CdCl<sub>2</sub> had a significant inhibitory effect on *Scenedesmus quadracauda* and that 0.061mg/L severely inhibited growth.

The EC50 value of 0.06mg/L Cd(II) as CdCl<sub>2</sub> reported by

Oyamada et al.<sup>33</sup> for *Chlorella vulgaris* is slightly higher than the EC50 values of 0.019, 0.005, 0.019, and 0.015 mg/L Cd(II) found in this investigation for *Ankistrodesmus*, *Scenedesmus*, *Selenastrum*, and *Microcoleus*, respectively. No marked difference in sensitivity to Cd(II) was noticed between blue-green and green algae.

The algistatic-algicidal values of 0.1-0.4 mg/L Hg(II) for the green algae and approximately 1.0mg/L Hg(II) for *Microcoleus* compare favorably with values reported in the literature. Mercuric chloride concentrations of 2.0 mg/L for *Chlamydomonas reinhardtii* y<sup>-1</sup> 0.037mg/L for *Chlorella vulgaris*, and 1.0mg/L for *Chlorella pyrenoidosa* were reported as algistatic-algicidal by Shen.<sup>34</sup> Tompkins & Blinn<sup>35</sup> reported algistatic-algicidal responses at 0.1mg/L and 0.5mg/L Hg(II) as HgCl<sub>2</sub> for *Fragilaria crotonensis* and *Asterionella formosa*, respectively, and Oh & Koh<sup>36</sup> reported an algistatic-algicidal response at 0.2mg/L Hg(II) as inorganic mercury for *Chlorella vulgaris*. Tezuka & Takasaki<sup>37</sup> reported less than 25µg/L Hg(II) as phenylmercuric acetate resulted in an algicidal-algistatic response in *Phaeodactylum tricornutum*, *Chlorella* sp. and *Chlamydomonas* sp. The EC50 values of 0.078, 0.085, 0.033, and 0.253 mg/L Hg(II) for *Ankistrodesmus*, *Scenedesmus*, *Selenastrum* and *Microcoleus* respectively, are smaller than the 1.03 mg/L Hg(II) as HgCl<sub>2</sub> reported by Patil<sup>38</sup> for *Chlorella vulgaris*. The blue-green alga *Microcoleus* appears to be somewhat less sensitive to Hg(II), but this could be due to differences in test media or to the fact that the *Microcoleus* was not axenic.

Little information is available concerning the toxicity of selenate to freshwater algae. Tomioka et al.<sup>39</sup> reported an algistatic-algicidal response of the blue-green alga *Anacystis nidulans* at 20mg/L Se(VI) as Na<sub>2</sub>SeO<sub>4</sub>. The algistatic-algicidal levels in this investigation ranged from 1.7mg/L Se(VI) for *Selenastrum* to >50 mg/L for *Microcoleus*. No response values for green algae exposed to Se(VI) were found in the literature. Although it appears that green algae are more sensitive to Se(VI) than are blue-green algae, the antagonistic response between sulfur and selenium in the different media complicates this relationship. Our EC50 values, which ranged from 0.033-8.511 mg/L Se(VI), are not directly comparable to any values in the literature.

Sun et al.<sup>40</sup> reported an algistatic-algicidal response of *Chlorella vulgaris* to 12mg/L Se(IV) as Na<sub>2</sub>SeO<sub>3</sub>. Kumar & Prakash<sup>41</sup> reported LD50 values for *Anabaena variabilis* and *Anacystis nidulans* at 13mg/L and 31mg/L Se(IV) as Na<sub>2</sub>SeO<sub>3</sub>, respectively, and at 18mg/L and 42mg/L Se(VI) as Na<sub>2</sub>SeO<sub>4</sub>, respectively. The response values obtained from the literature for Se(IV) are not, however, directly comparable to Se(VI) response values.

The demonstrated interspecies variation in sensitivity to the test substances in this study indicates that more than one algal species should be utilized when assessing the potential impact of toxic trace elements on an ecosystem. The use of a standard species such as *Selenastrum capricornutum* for interlaboratory comparisons is recommended for purposes of quality control. However, dependence on a single species can provide erroneous results concerning the potential toxicity of a toxic element in a given ecosystem. The highly toxic nature of potential pollutants from coal-fired generating plants to indigenous algae emphasizes the need for minimizing stack effluent pollutants and detaining scrubber ash slurry for proper disposal. Efforts should be made to maintain trace elements within concentration ranges which are compatible with the natural functioning of the associated aquatic ecosystems.

## Acknowledgements

The author is thankful to Prof. Loai Aljerf, the editor in the Biodiversity International Journal for his scientific assistance and his appreciated backup which eased reaching this paper to its final edition.

## Conflict of interest

Author declares that no competing interests exist.

## References

1. Spitz N, Saveliev R, Perelman M, et al. Firing a sub-bituminous coal in pulverized coal boilers configured for bituminous coals. *Fuel*. 2008;87(8-9):1534–1542.
2. Yinghui L, Chuguang Z, Quanhai W. Speciation of most volatile toxic trace elements during coal combustion. *Asia-Pacific Journal of Chemical Engineering*. 2008;11(3-4):381–394.
3. Qi LQ, Zhang YJ. Investigation on the characteristic of fly ash in electrostatic precipitators. *Advanced Materials Research*. 2012;518:869–872.
4. Mahlaba JS, Kearsley EP, Kruger RA. Physical, chemical and mineralogical characterisation of hydraulically disposed fine coal ash from SASOL Synfuels. *Fuel*. 2011;90(7):2491–2500.
5. Mwakabona HT. The growing global demand for element gallium: electrical and electronic waste and coal fly ash as alternative sources for its sustainable supply. *International Journal of Science and Research*. 2017;6(1):1287–1293.
6. Jiru M. Coal-fired power plants and their impact on ecosystems health. *Physical Science International Journal*. 2017;16(4):1–7.
7. Hao Y, Li Q, Pan Y, et al. Heavy metals distribution characteristics of FGD gypsum samples from Shanxi province 12 coal-fired power plants and its potential environmental impacts. *Fuel*. 2017;209:238–245.
8. Zamri SNAM, Sulaiman MA, Ismail S, et al. Bacto-Agar and commercial agar as binder for porous zinc anode. *Solid State Phenomena*. 2017;264:136–139.
9. Wang P, Li Z, Ismail S, et al. Optimization of microalgal bead preparation with *Scenedesmus obliquus* for both nutrient removal and lipid production. *Ecological Engineering*. 2016;92:236–242.
10. Nojiri H, Okoshi M. Crack suppression of silica glass formed by zoned F2 laser-induced photochemical surface modification of hard silicone thin film coating on polycarbonate. *Japanese Journal of Applied Physics*. 2016;55(12):122701.
11. Hai R, Feng H, Wang W. Adsorption of mercury(II) by beer yeast immobilized in chitosan/silicone leg. *Asian Journal of Chemistry*. 2013;25(12):6528–6530.
12. Lee DK, Cho YH, Lee JW, et al. Wettability of microstructured Pyrex glass with hydrophobic and hydrophilic properties. *Surface and Coatings Technology*. 2017;319:213–218.
13. Jackson G, Seymour LF. Formation constants at high ionic strength—II. The ionic strength correction of formation constants using a simplified pitzer equation. *Talanta*. 1995;42(1):9–16.
14. Eklund B. Review of the use of Ceramium tenuicorne growth inhibition test for testing toxicity of substances, effluents, products sediment and soil. *Estuarine, Coastal and Shelf Science*. 2017;195:88–97.
15. Awasthi M, Das DN. Heavy metal toxicity on nitrate reductase activity of free and immobilized algal cells. *International Journal on Algae*. 2004;6(2):151–157.
16. Tawarada K, Sasaki K, Ohmura N, et al. Preparation of anti-cadmium-EDTA complex monoclonal antibody and its binding specificity. *Bunseki Kagaku*. 2003;52(8):583–587.
17. Horvatić J, Palijan G, Lukavský J. Algal responses to nutrient additions in water of Nature Park Kopački Rit (Croatia) by Miniaturized Algal Growth Bioassay. *Algological Studies*. 2003;110(1):117–126.
18. Vigneault B, Campbell PGC. Uptake of cadmium by freshwater green algae: effects of pH and aquatic humic substances. *Journal of Phycology*. 2005;41(1):55–61.
19. Nouchpramool S, Sumitra T, Leenanuphont V. Characterization of airborne particulates in bangkok urban area by neutron activation analysis. *Biological Trace Element Research*. 1999;71(1):181–187.
20. Pyta H, Rogula-Kozłowska W. Determination of mercury in size-segregated ambient particulate matter using CVAAS. *Microchemical Journal*. 2016;124:76–81.
21. Martins RJE, Pardo R, Boaventura RAR. Cadmium(II) and zinc(II) adsorption by the aquatic moss *Fontinalis antipyretica*: effect of temperature, pH and water hardness. *Water Research*. 2004;38(3):693–699.
22. Hart BA, Bertram PE. A cadmium-binding protein in a cadmium tolerant strain of *Chlorella pyrenoidosa*. *Environmental and Experimental Botany*. 1980;20(2):175–180.
23. Wong SL, Beaver JL. Algal bioassays to determine toxicity of metal mixtures. *Hydrobiologia*. 1980;74(3):199–208.
24. Müller R, Heinze J. Stabilization of oxidized sulfur and selenium centers in organic chalcogens by three-electron sulfur-sulfur, selenium-selenium, and sulfur-selenium bonds. A theoretical study using the semiempirical PM3 method. *Phosphorus, Sulfur, and Silicon and the Related Elements*. 1998;141(1):111–134.
25. Finley HE, McLaughlin D. Cultivation of the Peritrich *Tetrotrochidium heneguyi* in axenic and non-axenic media. A theoretical study using the semiempirical PM3 method. *The Journal of Protozoology*. 1965;12(1):41–47.
26. Pearson TW. Effect of mercurial perturbation on avian uterine calcium transport. *Cell Calcium*. 1981;2(2):137–144.
27. Rijstenbil JW, Gerringa LJ. Interactions of algal ligands, metal complexation and availability, and cell responses of the diatom *Ditylum brightwellii* with a gradual increase in copper. *Aquatic Toxicology*. 2002;56(2):115–131.
28. Rainbow PS. Trace metal bioaccumulation: Models, metabolic availability and toxicity. *Environment International*. 2007;33(4):576–582.
29. Jiang X, Ren C, Hu C, et al. Isolation and algicidal characterization of *Bowmanella denitrificans* S088 against *Chlorella vulgaris*. *World Journal of Microbiology and Biotechnology*. 2013;30(2):621–629.
30. Mekhalif M, Amara S, Robert S, et al. Effect of environmental conditions on various enzyme activities and triacylglycerol contents in cultures of the freshwater diatom, *Asterionella formosa* (Bacillariophyceae). *Biochimie*. 2014;101:21–30.



31. Cheng J, Qiu H, Chang Z, et al. The effect of cadmium on the growth and antioxidant response for freshwater algae *Chlorella vulgaris*. *Springerplus*. 2016;5(1):1290.
32. Liu A, Tian D, Xiang Y, et al. Biochar improved growth of an important medicinal plant (*Salvia miltiorrhiza* Bunge) and inhibited its cadmium uptake. *Journal of Plant Biology & Soil Health*. 2016;3(2):6.
33. Oyamada N, Takahashi G, Ishizaki M. Methylation of inorganic selenium compounds by freshwater green Algae, *Ankistrodesmus* sp., *Chlorella vulgaris* and *Selenastrum* sp. *Eisei kagaku*. 1991;37(2):83–88.
34. Shen S. Genetic diversity analysis with ISSR PCR on green algae *Chlorella vulgaris* and *Chlorella pyrenoidosa*. *Chinese Journal of Oceanology and Limnology* 2008;26(4):380–384.
35. Tompkins T, Blinn DW. The effect of mercury on the growth rate of *Fragilaria crotonensis* Kitton and *Asterionella formosa* Hass. *Hydrobiologia*. 1976;49(2):111–116.
36. Oh S, Koh SC. Chlorophyll a fluorescence response to mercury stress in the freshwater Microalga *Chlorella vulgaris*. *Journal of Environmental Science International*. 2013;22(6):705–715.
37. Tezuka T, Takasaki Y. Biodegradation of phenylmercuric acetate by organomercury-resistant *Penicillium* sp. MR-2. *Agricultural and Biological Chemistry*. 2013;52(12):3183–3185.
38. Patil HS. The role of *Ankistrodesmus falcatus* and *Scenedesmus quadricauda* in sewage purification. *Bioresource Technology*. 1991;37(2):121–126.
39. Tomioka N, Shinozaki K, Sugiura M. Molecular cloning and characterization of ribosomal RNA genes from a blue-green alga, *Anacystis nidulans*. *MGG Molecular & General Genetics*. 1981;184(3):359–363.
40. Sun X, Zhong Y, Huang Z, et al. Selenium accumulation in unicellular green Alga *Chlorella vulgaris* and its effects on antioxidant enzymes and content of photosynthetic pigments. *PLoS ONE*. 2014;9(11):e112270.
41. Kumar HD, Prakash G. Toxicity of selenium to the Blue-green Algae, *Anacystis nidulans* and *Anabaena variabilis*. *Annals of Botany*. 1971;35(3):697–705.