

Satellite Communications

Lecture (3)


Chapter 2.1

This chapter is about how earth orbit is achieved, the laws that describe the motion of an object orbiting another body, how satellites maneuver in space, and the determination of the look angle to a satellite from the earth using ephemeris data that describe the orbital trajectory of the satellite.

To achieve a stable orbit around the earth, a spacecraft must first be beyond the bulk of the earth's atmosphere, i.e., in what is popularly called space. There are many definitions of space. U.S. astronauts are awarded their "space wings" if they fly at an altitude that exceeds 50 miles (~ 80 km); some international treaties hold that the space frontier above a given country begins at a height of 100 miles (~ 160 km).

To appreciate the basic laws that govern celestial mechanics, we will begin first with the fundamental Newtonian equations that describe the motion of a body. We will then give some coordinate axes within which the orbit of the satellite can be set and determine the various forces on the earth satellite.

Gravitational Force



- Newton's 2nd Law: $\vec{F} = m \cdot \vec{a}$

- Newton's Law Of Universal Gravitation (assuming point masses or spheres): $\vec{F}_{\text{body } 2} = \frac{Gm_1m_2}{r^2} \cdot \vec{u}_{2 \rightarrow 1}$

- Putting these together: $\vec{a}_{\text{body } 2} = \frac{Gm_1}{r^2} \cdot \vec{u}_{2 \rightarrow 1}$

2.1 ORBITAL MECHANICS

Developing the Equations of the Orbit

Newton's laws of motion can be encapsulated into four equations:

$$s = ut + \left(\frac{1}{2}\right)at^2 \quad (2.1a)$$

$$v^2 = u^2 + 2at \quad (2.1b)$$

$$v = u + at \quad (2.1c)$$

$$P = ma \quad (2.1d)$$

where s is the distance traveled from time $t = 0$; u is the initial velocity of the object at time $t = 0$ and v the final velocity of the object at time t ; a is the acceleration of the object; P is the force acting on the object; and m is the mass of the object. Note that the acceleration can be positive or negative, depending on the direction it is acting with respect to the velocity vector. Of these four equations, it is the last one that helps us understand the motion of a satellite in a stable orbit neglecting any drag or other perturbing forces). Put into words, Eq. (2.1d) states that the force acting on a body is equal to the mass of the body multiplied by the resulting acceleration of the body.

the lighter the mass of the body, the higher the acceleration

Kinematics & Newton's Law

s = Distance traveled in time, t


u = Initial Velocity at $t = 0$

v = Final Velocity at time = t

a = Acceleration

F = Force acting on the object

Newton's
Second Law



- $s = ut + (1/2)at^2$
- $v^2 = u^2 + 2at$
- $v = u + at$
- $F = ma$

there are two main forces acting on a satellite:

: a centrifugal force

the kinetic energy of the satellite, which attempts to fling the satellite into a higher orbit

a centripetal force

due to the gravitational attraction of the planet about which the satellite is orbiting, which attempts to pull the satellite down toward the planet.

If these two forces are equal, the satellite will remain in a stable orbit.


Force = mass $\times$ acceleration and the unit of force is a Newton, with the notation N . A Newton is the force required to accelerate a mass of 1 kg with an acceleration of 1 m/s². The underlying units of a Newton are therefore (kg) $\times$ m/s². In Imperial Units one Newton = 0.2248 ft lb. The standard acceleration due to gravity at the earth's surface is 9.80665×10^{-3} km/s², which is often quoted as 981 cm/s². This value decreases

FORCE ON A SATELLITE : 1

- **Force = Mass $\times$ Acceleration**
- Unit of Force is a (***Newton***)
- A ***Newton*** is the force required to accelerate 1 kg by 1 m/s²
- Underlying units of a ***Newton*** are therefore (kg) $\times$ (m/s²)
- In Imperial Units 1 ***Newton*** = 0.2248 ft lb.

ACCELERATION FORMULA

- $a = \text{acceleration due to gravity} = \mu / r^2 \text{ km/s}^2$
- $r = \text{radius from center of earth}$
- $\mu = \text{universal gravitational constant}$
 G multiplied by the mass of the earth M_E
- μ is Kepler's constant and = $3.9861352 \times 10^5 \text{ km}^3/\text{s}^2$
- $G = 6.672 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$ or $6.672 \times 10^{-20} \text{ km}^3/\text{kg s}^2$ in the older units



The acceleration (a) due to gravity at a distance r from the earth is $a = \mu/r^2 \text{ km/s}^2$

μ is the product of the universal gravitational constant G and the mass of the earth M_E , this is kepler's constant and has the value $\mu = 3.986004418 \times 10^5 \text{ km}^3/\text{kg}$.

The universal gravitational constant is $G = 6.672 \times 10^{-11} \text{ Nm}^2/\text{kg}^2 = G = 6.672 \times 10^{-20} (\text{km}^3/\text{kg s}^2)$ in the older units.

Force = mass $\times$ acceleration

$$F_{IN} = m \times (\mu/r^2)$$

Centripetal force acting on satellite, $F_{IN} = m \times (GM_E/r^2)$

The centrifugal acceleration, $a = v^2/r$

The centrifugal force, $F_{OUT} = m \times (v^2/r)$

If the force on the satellite are balanced

$$F_{IN} = F_{OUT}$$

$$m \times \mu/r^2 = m \times v^2/r$$

The velocity v of a satellite in a circular orbit is,

$$v = (\mu/r)^{1/2}$$

The period of satellite's orbit, T

if the orbit is circular, the distance traveled by a satellite in one orbit around a planet is $2\pi r$, where r is the radius of the orbit from the satellite to the center of the planet. Since distance divided by velocity equals time to travel that distance, the period of the satellite's orbit, T , will be

$$T = (2\pi r)/v = (2\pi r)/[(\mu/r)^{1/2}] \quad (2.6)$$

$$T = (2\pi r^{3/2})/(\mu^{1/2})$$

TABLE 2.1 Orbital Velocity, Height, and Period of Four Satellite Systems

Satellite system	Orbital height (km)	Orbital velocity (km/s)	Orbital period (h min s)		
Intelsat (GEO)	35,786.03	3.0747	23	56	4.1
New-ICO (MEO)	10,255	4.8954	5	55	48.4
Skybridge (LEO)	1,469	7.1272	1	55	17.8
Iridium (LEO)	780	7.4624	1	40	27.0

Mean earth radius is 6378.137 km and GEO radius from the center of the earth is 42,164.17 km.

a geocentric coordinate system.




FIGURE 2.2 The initial coordinate system that could be used to describe the relationship between the earth and a satellite. A Cartesian coordinate system with the geographical axes of the earth as the principal axes is the simplest coordinate system to set up. The rotational axis of the earth is about the axis cz , where c is the center of the earth and cz passes through the geographic north pole. Axes cx , cy , and cz are mutually orthogonal axes, with cx and cy passing through the earth's geographic equator. The vector r locates the moving satellite with respect to the center of the earth.

the satellite mass m located at a vector distance $\mathbf{r}$ from the center of the earth.
the gravitational force $\mathbf{F}$ on the satellite is given by

$$\mathbf{F} = -\frac{GM_E m \mathbf{r}}{r^3}$$

M_E is the mass of the earth and $G = 6.672 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$.

But force = mass $\times$ acceleration

$$\mathbf{F} = m \frac{d^2 \mathbf{r}}{dt^2}$$

$$-\frac{\mathbf{r}}{r^3} \mu = \frac{d^2 \mathbf{r}}{dt^2}$$

Which yields

$$\frac{d^2 \mathbf{r}}{dt^2} + \frac{\mathbf{r}}{r^3} \mu = 0$$

(2.10)

This is a second-order linear differential equation and its solution will involve six undetermined constants called the *orbital elements*. The orbit described by these orbital elements can be shown to lie in a plane and to have a constant angular momentum. The solution to Eq. (2.10) is difficult since the second derivative of $\mathbf{r}$ involves the second derivative of the unit vector $\mathbf{r}$. To remove this dependence, a different set of coordinates can

be chosen to describe the location of the satellite such that the unit vectors in the three axes are constant. This coordinate system uses the plane of the satellite's orbit as the reference plane. This is shown in Figure 2.3.




FIGURE 2.3 The orbital plane coordinate system. In this coordinate system, the orbital plane of the satellite is used as the reference plane. The orthogonal axes x_0 and y_0 lie in the orbital plane. The third axis, z_0 , is perpendicular to the orbital plane. The geographical z -axis of the earth (which passes through the true North Pole and the center of the earth, c) does not lie in the same direction as the z_0 axis except for satellite orbits that are exactly in the plane of the geographical equator.

Expressing Eq. (2.10) in terms of the new coordinate axes x_0 , y_0 , and z_0 gives

$$\hat{x}_0 \left(\frac{d^2 x_0}{dt^2} \right) + \hat{y}_0 \left(\frac{d^2 y_0}{dt^2} \right) + \frac{\mu(x_0 \hat{x}_0 + y_0 \hat{y}_0)}{(x_0^2 + y_0^2)^{3/2}} = 0 \quad (2.11)$$

Equation (2.11) is easier to solve if it is expressed in a polar coordinate system rather than a Cartesian coordinate system. The polar coordinate system is shown in Figure 2.4.




FIGURE 2.4 Polar coordinate system in the plane of the satellite's orbit. The plane of the orbit coincides with the plane of the paper. The axis z_0 is straight out of the paper from the center of the earth, and is normal to the plane of the satellite's orbit. The satellite's position is described in terms of the radius from the center of the earth r_0 and the angle this radius makes with the x_0 axis, ϕ_0 .

With the polar coordinate system shown in Figure 2.4 and using the transformations

$$x_0 = r_0 \cos \phi_0 \quad (2.12a)$$

$$y_0 = r_0 \sin \phi_0 \quad (2.12b)$$

$$\hat{x}_0 = \hat{r}_0 \cos \phi_0 - \hat{\phi}_0 \sin \phi_0 \quad (2.12c)$$

$$\hat{y}_0 = \hat{\phi}_0 \cos \phi_0 + \hat{r}_0 \sin \phi_0 \quad (2.12d)$$

and equating the vector components of r_0 and ϕ_0 in turn in Eq. (2.11) yields

$$\frac{d^2 r_0}{dt^2} - r_0 \left(\frac{d\phi_0}{dt} \right)^2 = -\frac{\mu}{r_0^2} \quad (2.13)$$

and

$$r_0 \left(\frac{d^2 \phi_0}{dt^2} \right) + 2 \left(\frac{dr_0}{dt} \right) \left(\frac{d\phi_0}{dt} \right) = 0 \quad (2.14)$$

Using standard mathematical procedures, we can develop an equation for the radius of the satellite's orbit, r_0 , namely

$$r_0 = \frac{p}{1 + e \cos(\phi_0 - \theta_0)} \quad (2.15)$$

Where θ_0 is a constant and e is the eccentricity of an ellipse whose semilatus rectum p is given by

$$p = (h^2)/\mu \quad (2.16)$$

and h is the magnitude of the orbital angular momentum of the satellite. That the equation of the orbit is an ellipse is Kepler's first law of planetary motion.

Kepler's Three Laws


First Law: The orbit of each planet is an ellipse, with the Sun at a focus (1609).

Second Law: The line joining the planet to the Sun sweeps out equal areas in equal times (1609).

Third Law: The square of the period of a planet is proportional to the cube of its mean distance from the Sun (1619).

PHYSICAL LAWS


Kepler's 1st Law: Law of Ellipses



The orbits of the planets are ellipses with the sun at one focus


PHYSICAL LAWS

Kepler's 2nd Law: Law of Equal Areas
The line joining the planet to the center of the sun sweeps out equal areas in equal times



PHYSICAL LAWS

Kepler's 2nd Law: Law of Equal Areas



$$t_1 - t_0 = t_3 - t_2$$

$$\text{Area 1} = \text{Area 2}$$


Satellite travels at varying speeds

PHYSICAL LAWS

Kepler's 3rd Law: Law of Harmonics
 The squares of the periods of
 two planets' orbits are
 proportional to each other as
 the cubes of their semi-
 major axes:

$$T_1^2/T_2^2 = a_1^3/a_2^3$$

In English:
 Orbits with the same semi-
 major axis will have the
 same period




Kepler's Three Laws of Planetary Motion

Johannes Kepler (1571–1630) was a German astronomer and scientist who developed his three laws of planetary motion by careful observations of the behavior of the planets in the solar system over many years, with help from some detailed planetary observations by the Hungarian astronomer Tycho Brahe. Kepler's three laws are

1. The orbit of any smaller body about a larger body is always an ellipse, with the center of mass of the larger body as one of the two foci.
2. The orbit of the smaller body sweeps out equal areas in equal time (see Figure 2.5).
3. The square of the period of revolution of the smaller body about the larger body equals a constant multiplied by the third power of the semimajor axis of the orbital ellipse. That is, $T^2 = (4\pi^2 a^3)/\mu$ where T is the orbital period, a is the semimajor axis of the orbital ellipse, and μ is Kepler's constant. If the orbit is circular, then a becomes distance r , defined as before, and we have Eq. (2.6).

Describing the orbit of a satellite enables us to develop Kepler's second two laws.



Describing the Orbit of a Satellite

The quantity θ_0 in Eq. (2.15) serves to orient the ellipse with respect to the orbital plane axes x_0 and y_0 . Now that we know that the orbit is an ellipse, we can always choose x_0 and y_0 so that θ_0 is zero. We will assume that this has been done for the rest of this discussion. This now gives the equation of the orbit as

$$r_0 = \frac{p}{1 + e \cos \phi_0} \quad (2.17)$$

The path of the satellite in the orbital plane is shown in Figure 2.6. The lengths a and b of the semimajor and semiminor axes are given by

$$a = p/(1 - e^2) \quad (2.18)$$

$$b = a(1 - e^2)^{1/2} \quad (2.19)$$

The point in the orbit where the satellite is closest to the earth is called the *perigee* and the point where the satellite is farthest from the earth is called the *apogee*. The perigee and apogee are **always** exactly opposite each other. To make θ_0 equal to zero, we have chosen the x_0 axis so that both the apogee and the perigee lie along it and the x_0 axis is therefore the major axis of the ellipse.

The differential area swept out by the vector $\mathbf{r}_0$ from the origin to the satellite in time dt is given by

$$dA = 0.5r_0^2 \left(\frac{d\phi_0}{dt} \right) dt = 0.5h dt \quad (2.20)$$

Remembering that h is the magnitude of the orbital angular momentum of the satellite, the radius vector of the satellite can be seen to sweep out equal areas in equal times. This is Kepler's second law of planetary motion. By equating the area of the ellipse (πab) to the area swept out in one orbital revolution, we can derive an expression for the orbital period T as

$$T^2 = (4\pi^2 a^3)/\mu \quad (2.21)$$

$$T^2 = (4\pi^2 a^3)/\mu$$

$$T = (2\pi r^{3/2})/(\mu^{1/2})$$

This equation is the mathematical expression of Kepler's third law of planetary motion: the square of the period of revolution is proportional to the cube of the semimajor axis. (Note that this is the square of Eq. (2.6) and that in Eq. (2.6) the orbit was assumed to be circular such that semimajor axis a = semiminor axis b = circular orbit radius from the center of the earth r .) *Kepler's third law extends the result from Eq. (2.6), which was derived for a circular orbit, to the more general case of an elliptical orbit.* Equation (2.21) is extremely important in satellite communications systems. This equation determines the period of the orbit of any satellite, and it is used in every GPS receiver in the calculation of the positions of GPS satellites. Equation (2.21) is also used to find the orbital radius of a GEO satellite, for which the period T must be made exactly equal to the period of one revolution of the earth for the satellite to remain stationary over a point on the equator.

An important point to remember is that the period of revolution, T , is referenced to inertial space, namely, to the galactic background. The orbital period is the time the orbiting body takes to return to the same reference point in space with respect to the galactic background. Nearly always, the primary body will also be rotating and so the period of revolution of the satellite may be different from that perceived by an observer who is standing still on the surface of the primary body. This is most obvious with a geostationary earth orbit (GEO) satellite (see Table 2.1). The orbital period of a GEO satellite is exactly equal to the period of rotation of the earth, 23 h 56 min 4.1 s, but, to an observer on the ground, the satellite appears to have an infinite orbital period: it always stays in the same place in the sky.


geostationary vs. *geosynchronous* orbit.

To be perfectly geostationary, the orbit of a satellite needs to have three features: (a) it must be exactly circular (i.e., have an eccentricity of zero); (b) it must be at the correct altitude (i.e., have the correct period); and (c) it must be in the plane of the equator (i.e., have a zero inclination with respect to the equator). If the inclination of the satellite is not zero and/or if the eccentricity is not zero, but the orbital period is correct, then the satellite will be in a *geosynchronous* orbit. The position of a geosynchronous satellite will appear to oscillate about a mean look angle in the sky with respect to a stationary observer on the earth's surface. The orbital period of a GEO satellite, 23 h 56 min 4.1 s, is one sidereal day. A sidereal day is the time between consecutive crossings of any particular longitude on the earth by any star, other than the sun. The mean solar day of 24 h is the time between any consecutive crossings of any particular longitude by the sun, and is the time between successive sunrises (or sunsets) observed at one location on earth, averaged over an entire year. Because the earth moves round the sun once per 365 $\frac{1}{4}$ days, the solar day is $1440/365.25 = 3.94$ min longer than a sidereal day.

Solar vs. Sidereal Day


- A ***sidereal day*** is the time between consecutive crossings of any particular longitude on the earth by any star other than the sun.
- A ***solar day*** is the time between consecutive crossings of any particular longitude of the earth by the sun-earth axis.
 - Solar day = EXACTLY 24 hrs
 - Sidereal day = 23 h 56 min. 4.091 s
- Why the difference?
 - By the time the Earth completes a full rotation with respect to an external point (not the sun), it has already moved its center position with respect to the sun. The extra time it takes to cross the sun-earth axis, averaged over 4 full years (because every 4 years one has 366 days) is of about 3.93 minutes per day.

Calculation next page



LOCATING THE SATELLITE IN ORBIT: 1

Start with Fig. 2.6 in Text



NOTE: **Perigee** and **Apogee** are on opposite sides of the orbit

31




FIGURE 2.6 The orbit as it appears in the orbital plane. The point O is the center of the earth and the point C is the center of the ellipse. The two centers do not coincide unless the eccentricity, e , of the ellipse is zero (i.e., the ellipse becomes a circle and $a = b$). The dimensions a and b are the semimajor and semiminor axes of the orbital ellipse, respectively.

Locating the Satellite in the Orbit

Consider now the problem of locating the satellite in its orbit. The equation of the orbit may be rewritten by combining Eqs. (2.15) and (2.18) to obtain

$$r_0 = \frac{a(1 - e^2)}{1 + e \cos \phi_0} \quad (2.22)$$

The angle ϕ_0 (see Figure 2.6) is measured from the x_0 axis and is called the *true anomaly*. [*Anomaly* was a measure used by astronomers to mean a planet's angular distance from its perihelion (closest approach to the sun), measured as if viewed from the sun. The term was adopted in celestial mechanics for all orbiting bodies.] Since we defined the positive x_0 axis so that it passes through the perigee, ϕ_0 measures the angle from the perigee to the instantaneous position of the satellite. The rectangular coordinates of the satellite are given by

$$x_0 = r_0 \cos \phi_0 \quad (2.23)$$


$$y_0 = r_0 \sin \phi_0 \quad (2.24)$$

As noted earlier, the orbital period T is the time for the satellite to complete a revolution in inertial space, traveling a total of 2π radians. The average angular velocity η is thus

$$\eta = (2\pi)/T = (\mu^{1/2})/(a^{3/2}) \quad (2.25)$$

If the orbit is an ellipse, the instantaneous angular velocity will vary with the position of the satellite around the orbit. If we enclose the elliptical orbit with a *circumscribed circle* of radius a (see Figure 2.7), then an object going around the circumscribed circle with a constant angular velocity η would complete one revolution in exactly the same period T as the satellite requires to complete one (elliptical) orbital revolution.

Consider the geometry of the circumscribed circle as shown in Figure 2.7. Locate the point (indicated as A) where a vertical line drawn through the position of the satellite intersects the circumscribed circle. A line from the center of the ellipse (C) to this point (A) makes an angle E with the x_0 axis; E is called the *eccentric anomaly* of the satellite.



It is related to the radius r_0 by

$$r_0 = a(1 - e \cos E) \quad (2.26)$$

Thus

$$a - r_0 = ae \cos E \quad (2.27)$$

We can also develop an expression that relates eccentric anomaly E to the average angular velocity η , which yields

$$\eta dt = (1 - e \cos E) dE \quad (2.28)$$

Let t_p be the time of perigee. This is simultaneously the time of closest approach to the earth; the time when the satellite is crossing the x_o axis; and the time when E is zero. If we integrate both sides of Eq. (2.28), we obtain

$$\eta(t - t_p) = E - e \sin E \quad (2.29)$$

The left side of Eq. (2.29) is called the mean anomaly, M . Thus

$$M = \eta(t - t_p) = E - e \sin E \quad (2.30)$$

The mean anomaly M is the arc length (in radians) that the satellite would have traversed since the perigee passage if it were moving on the circumscribed circle at the mean angular velocity η .

If we know the time of perigee, t_p , the eccentricity, e , and the length of the semi-major axis, a , we now have the necessary equations to determine the coordinates (r_0, ϕ_0)

locating the satellite at the point (x_0, y_0, z_0)

and (x_0, y_0) of the satellite in the orbital plane. The process is as follows

1. Calculate η using Eq. (2.25). $\eta = (2\pi)/T = (\mu^{1/2})/(a^{3/2})$
2. Calculate M using Eq. (2.30). $M = \eta(t - t_p) = E - e \sin E$
3. Solve Eq. (2.30) for E .
4. Find r_0 from E using Eq. (2.27). $a - r_0 = ae \cos E$

$$r_0 = \frac{a(1 - e^2)}{1 + e \cos \phi_0}$$
5. Solve Eq. (2.22) for ϕ_0 .
6. Use Eqs. (2.23) and (2.24) to calculate x_0 and y_0 . $x_0 = r_0 \cos \phi_0$
 $y_0 = r_0 \sin \phi_0$

Now we must locate the orbital plane with respect to the earth.

Locating the Satellite with Respect to the Earth

At the end of the last section, we summarized the process for locating the satellite at the point (x_0, y_0, z_0) in the rectangular coordinate system of the orbital plane. The location was with respect to the center of the earth. In most cases, we need to know where the satellite is from an observation point that is not at the center of the earth. We will therefore develop the transformations that permit the satellite to be located from a point on the rotating surface of the earth. We will begin with a geocentric equatorial coordinate system as shown in Figure 2.8. The rotational axis of the earth is the z_i axis, which is through the geographic North Pole. The x_i axis is from the center of the earth toward a fixed location in space called the *first point of Aries* (see Figure 2.8). This coordinate system moves through space; it translates as the earth moves in its orbit around the sun, but it does not rotate as the earth rotates. The x_i direction is always the same, whatever the earth's position around the sun and is in the direction of the first point of Aries. The (x_i, y_i) plane contains the earth's equator and is called the equatorial plane.

Locating the Satellite with Respect to the Earth





FIGURE 2.8 The geocentric equatorial system. This geocentric system differs from that shown in Figure 2.1 only in that the x_i axis points to the first point of Aries. The first point of Aries is the direction of a line from the center of the earth through the center of the sun at the vernal equinox (about March 21 in the Northern Hemisphere), the instant when the subsolar point crosses the equator from south to north. In the above system, an object may be located by its right ascension RA and its declination δ .

Angular distance measured eastward in the equatorial plane from the x_i axis is called **right ascension** and given the symbol RA . The two points at which the orbit penetrates the equatorial plane are called **nodes**; the satellite moves upward through the equatorial plane at the **ascending node** and downward through the equatorial plane at the **descending node** given the conventional picture of the earth, with north at the top, which is in the direction of the positive z axis for the earth centered coordinate set. Remember that in space there is no *up* or *down*; that is a concept we are familiar with because of gravity at the earth's surface. For a weightless body in space, such as an orbiting spacecraft, up and down have no meaning unless they are defined with respect to a reference point. **The right ascension of the ascending node** is called Ω . The angle that the orbital plane makes with the equatorial plane (the planes intersect at the line joining the nodes) is called the **inclination, i** . Figure 2.9 illustrates these quantities.



The variables Ω and i together locate the orbital plane with respect to the equatorial plane. To locate the orbital coordinate system with respect to the equatorial coordinate system we need ω , the *argument of perigee west*. This is the angle measured along the orbit from the ascending node to the perigee.

Standard time for space operations and most other scientific and engineering purposes is *universal time (UT)*, also known as *zulu time (Z)*. This is essentially the mean solar time at the Greenwich Observatory near London, England. Universal time is measured in hours, minutes, and seconds or in fractions of a day. It is 5 h later than Eastern Standard Time, so that 07:00 EST is 12:00:00 h UT. The civil or calendar day begins at 00:00:00 hours UT, frequently written as 0 h. This is, of course, midnight (24:00:00) on the previous day. Astronomers employ a second dating system involving *Julian days* and *Julian dates*. Julian days start at noon UT in a counting system whereby noon on December 31, 1899, was the beginning of Julian day 2415020, usually written 241 5020. These are extensively tabulated in reference 2 and additional information is in reference 14. As an example, noon on December 31, 2000, the eve of the twenty-first century, is the start of Julian day 245 1909. Julian dates can be used to indicate time by appending a decimal fraction; 00:00:00 h UT on January 1, 2001—zero hour, minute, and second for the third millennium A.D.—is given by Julian date 245 1909.5. To find the exact position of an orbiting satellite at a given instant in time requires knowledge of the orbital elements.

Orbital Elements

To specify the absolute (i.e., the inertial) coordinates of a satellite at time t , we need to know six quantities. (This was evident earlier when we determined that a satellite's equation of motion was a second order vector linear differential equation.) These quantities are called the orbital elements. More than six quantities can be used to describe a unique orbital path and there is some arbitrariness in exactly which six quantities are used. We have chosen to adopt a set that is commonly used in satellite communications: eccentricity (e), semimajor axis (a), time of perigee (t_p), right ascension of ascending node (Ω), inclination (i), and argument of perigee (ω). Frequently, the mean anomaly (M) at a given time is substituted for t_p .

EXAMPLE 2.1.1 Geostationary Satellite Orbit Radius

The earth rotates once per sidereal day of 23 h 56 min 4.09 s. Use Eq. (2.21) to show that the radius of the GEO is 42,164.17 km as given in Table 2.1.

Answer Equation (2.21) gives the square of the orbital period in seconds

$$T^2 = (4\pi^2 a^3)/\mu$$

Rearranging the equation, the orbital radius a is given by

$$a^3 = T^2 \mu / (4\pi^2)$$

For one sidereal day, $T = 86,164.09$ s. Hence

$$a^3 = (86,164.1)^2 \times 3.986004418 \times 10^5 / (4\pi^2) = 7.496020251 \times 10^{13} \text{ km}^3$$

$$a = 42,164.17 \text{ km}$$

This is the orbital radius for a geostationary satellite, as given in Table 2.1. ■

EXAMPLE 2.1.2 Low Earth Orbit

The Space Shuttle is an example of a low earth orbit satellite. Sometimes, it orbits at an altitude of 250 km above the earth's surface, where there is still a finite number of molecules from the atmosphere. The mean earth's radius is approximately 6378.14 km. Using these figures, calculate the period of the shuttle orbit when the altitude is 250 km and the orbit is circular. Find also the linear velocity of the shuttle along its orbit.

Answer The radius of the 250-km altitude Space Shuttle orbit is $(r_e + h) = 6378.14 + 250.0 = 6628.14$ km

From Eq. 2.21, the period of the orbit is T where

$$\begin{aligned} T^2 &= (4\pi^2 a^3)/\mu = 4\pi^2 \times (6628.14)^3 / 3.986004418 \times 10^5 \text{ s}^2 \\ &= 2.88401145 \times 10^7 \text{ s}^2 \end{aligned}$$

Hence the period of the orbit is

$$T = 5370.30 \text{ s} = 89 \text{ min } 30.3 \text{ s.}$$

This orbit period is about as small as possible. At a lower altitude, friction with the earth's atmosphere will quickly slow the Shuttle down and it will return to earth. Thus, all spacecraft in stable earth orbit have orbital periods exceeding 89 min 30 s.

The circumference of the orbit is $2\pi a = 41,645.83$ km.

Hence the velocity of the Shuttle in orbit is

$$2\pi a/T = 41,645.83/5370.13 = 7.755 \text{ km/s}$$

Alternatively, you could use Eq. (2.5): $v = (\mu/r)^{1/2}$. The term $\mu = 3.986004418 \times 10^5 \text{ km}^3/\text{s}^2$ and the term $r = (6378.14 + 250.0) \text{ km}$, yielding $v = 7.755 \text{ km/s}$.

Note: If μ and r had been quoted in units of m^3/s^2 and m , respectively, the answer would have been in meters/second. Be sure to keep the units the same during a calculation procedure.

A velocity of about 7.8 km/s is a typical velocity for a low earth orbit satellite. As the altitude of a satellite increases, its velocity becomes smaller. ■

EXAMPLE 2.1.3 Elliptical orbit

A satellite is in an elliptical orbit with a perigee of 1000 km and an apogee of 4000 km. Using a mean earth radius of 6378.14 km, find the period of the orbit in hours, minutes, and seconds, and the eccentricity of the orbit.

Answer The major axis of the elliptical orbit is a straight line between the apogee and perigee, as seen in Figure 2.7. Hence, for a semimajor axis length a , earth radius r_e , perigee height h_p , and apogee height h_a ,

$$2a = 2r_e + h_p + h_a = 2 \times 6378.14 + 1000.0 + 4000.0 = 17,756.28 \text{ km}$$

Thus the semimajor axis of the orbit has a length $a = 8878.14$ km. Using this value of a in Eq. (2.21) gives an orbital period T seconds where



$$\begin{aligned} T^2 &= (4\pi^2 a^3)/\mu = 4\pi^2 \times (8878.14)^3 / 3.986004418 \times 10^5 \text{ s}^2 \\ &= 6.930872802 \times 10^7 \text{ s}^2 \\ T &= 8325.1864 \text{ s} = 138 \text{ min } 45.19 \text{ s} = 2 \text{ h } 18 \text{ min } 45.19 \text{ s} \end{aligned}$$



The eccentricity of the orbit is given by e , which can be found from Eq. (2.27) by considering the instant at which the satellite is at perigee. Referring to Figure 2.7, when the satellite is at perigee, the eccentric anomaly $E = 0$ and $r_0 = r_e + h_p$. From Eq. (2.27), at perigee

$$r_0 = a(1 - e \cos E) \quad \text{and} \quad \cos E = 1$$

Hence

$$\begin{aligned} r_e + h_p &= a(1 - e) \\ e &= 1 - (r_e + h_p)/a = 1 - 7,378.14/8878.14 = 0.169 \end{aligned}$$






- Book Lecture notes

LOCATING THE SATELLITE IN ORBIT: 1

Start with Fig. 2.6 in Text



NOTE: **Perigee** and **Apogee** are on opposite sides of the orbit

LOCATING THE SATELLITE IN ORBIT: 2


- Need to develop a procedure that will allow the average angular velocity to be used
- If the orbit is not circular, the procedure is to use a ***Circumscribed Circle***
- A circumscribed circle is a circle that has a radius equal to the semi-major axis length of the ellipse and also has the same center

See next slide

53

LOCATING THE SATELLITE IN ORBIT: 3

Fig. 2.7 in the text



54

ORBIT CHARACTERISTICS

Semi-Axis Lengths of the Orbit

$$a = \frac{p}{1 - e^2} \quad \text{where} \quad p = \frac{h^2}{\mu}$$

See eq. (2.18)
and (2.16)

and h is the magnitude of
the angular momentum

$$b = a(1 - e^2)^{1/2} \quad \text{where} \quad e = \frac{h^2 C}{\mu}$$

See eqn.
(2.19)

and e is the eccentricity of the orbit

55

ORBIT ECCENTRICITY

- If a = semi-major axis,
 b = semi-minor axis, and
 e = eccentricity of the orbit ellipse,
then

$$e = \frac{a - b}{a + b}$$

NOTE: For a circular orbit, $a = b$ and $e = 0$

56

Time reference:

- t_p Time of Perigee = Time of closest approach to the earth, at the same time, time the satellite is crossing the x_0 axis, according to the reference used.
- $t - t_p$ = time elapsed since satellite last passed the perigee.

57

ORBIT DETERMINATION 1: Procedure:

Given the time of perigee t_p , the eccentricity e and the length of the semimajor axis a :

- η Average Angular Velocity (eqn. 2.25)
- M Mean Anomaly (eqn. 2.30)
- E Eccentric Anomaly (solve eqn. 2.30)
- r_0 Radius from orbit center (eqn. 2.27)
- ϕ_0 True Anomaly (solve eq. 2.22)
- x_0 and y_0 (using eqn. 2.23 and 2.24)

58

ORBIT DETERMINATION 2:

- ***Orbital Constants*** allow you to determine coordinates (r_o, φ_o) and (x_o, y_o) in the orbital plane
- Now need to locate the orbital plane with respect to the earth
- More specifically: need to locate the orbital location with respect to a **point on the surface of the earth**

59

LOCATING THE SATELLITE WITH RESPECT TO THE EARTH

- The orbital constants define the orbit of the satellite with respect to the **CENTER** of the earth
- To know where to look for the satellite in space, we must relate the orbital plane and time of perigee to the earth's axis

NOTE: Need a ***Time Reference*** to locate the satellite. The time reference most often used is the ***Time of Perigee, t_p***

60

GEOCENTRIC EQUATORIAL COORDINATES - 1

- z_i axis Earth's rotational axis (N-S poles with N as positive z)
- x_i axis In equatorial plane towards **FIRST POINT OF ARIES**
- y_i axis Orthogonal to z_i and x_i

NOTE: The *First Point of Aries* is a line from the center of the earth through the center of the sun at the vernal equinox (spring) in the northern hemisphere


61

GEOCENTRIC EQUATORIAL COORDINATES - 2

Fig. 2.8 in text

RA = Right Ascension
(in the x_i, y_i plane)

δ = Declination (the angle from the x_i, y_i plane to the satellite radius)



NOTE: Direction to *First Point of Aries* does **NOT** rotate with earth's motion around; the direction only *translates*

62


LOCATING THE SATELLITE - 1

- Find the ***Ascending Node***
 - Point where the satellite crosses the equatorial plane from South to North
 - Define Ω and i Inclination
 - Define ω Right Ascension of the Ascending Node (= ***RA*** from Fig. 2.6 in text)
- See next slide

63


DEFINING PARAMETERS

Fig. 2.9 in text



64

DEFINING PARAMETERS 2



(Source: M.Richaria, Satellite Communication Systems, Fig.2.9) 65

LOCATING THE SATELLITE - 2

- Ω and i together locate the Orbital plane with respect to the Equatorial plane.
- ω locates the Orbital coordinate system with respect to the Equatorial coordinate system.

66

LOCATING THE SATELLITE - 2

- Astronomers use **Julian Days** or **Julian Dates**
- Space Operations are in ***Universal Time Constant*** (UTC) taken from Greenwich Meridian (This time is sometimes referred to as ***“Zulu”***)
- To find exact position of an orbiting satellite at a given instant, we need the ***Orbital Elements***

67

ORBITAL ELEMENTS (P. 29)

- Ω Right Ascension of the Ascending Node
- i Inclination of the orbit
- ω Argument of Perigee (See Figures 2.6 & 2.7 in the text)
- t_p Time of Perigee
- e Eccentricity of the elliptical orbit
- a Semi-major axis of the orbit ellipse (See Fig. 2.4 in the text)

68

conclusions

Definition of terms for earth-orbiting satellite

- **Apogee** The point farthest from earth. Apogee height is shown as h_a in Fig
- **Perigee** The point of closest approach to earth. The perigee height is shown as h_p
- **Line of apsides** The line joining the perigee and apogee through the center of the earth.
- **Ascending node** The point where the orbit crosses the equatorial plane going from south to north.
- **Descending node** The point where the orbit crosses the equatorial plane going from north to south.
- **Line of nodes** The line joining the ascending and descending nodes through the center of the earth.
- **Inclination** The angle between the orbital plane and the earth's equatorial plane. It is measured at the ascending node from the equator to the orbit, going from east to north. The inclination is shown as i in Fig.
- **Mean anomaly** M gives an average value of the angular position of the satellite with reference to the perigee.
- **True anomaly** is the angle from perigee to the satellite position, measured at the earth's center. This gives the true angular position of the satellite in the orbit as a function of time.




Figure 2.3 Apogee height h_a , perigee height h_p , and inclination i . l_a is the line of apsides.

Definition of terms for earth-orbiting satellite

- **Prograde orbit** An orbit in which the satellite moves in the same direction as the earth's rotation. The inclination of a prograde orbit always lies between 0 and 90°.
- **Retrograde orbit** An orbit in which the satellite moves in a direction counter to the earth's rotation. The inclination of a retrograde orbit always lies between 90 and 180°.
- **Argument of perigee** The angle from ascending node to perigee, measured in the orbital plane at the earth's center, in the direction of satellite motion.
- **Right ascension of the ascending node** To define completely the position of the orbit in space, the position of the ascending node is specified. However, because the earth spins, while the orbital plane remains stationary the longitude of the ascending node is not fixed, and it cannot be used as an absolute reference. For the practical determination of an orbit, the longitude and time of crossing of the ascending node are frequently used. However, for an absolute measurement, a fixed reference in space is required. The reference chosen is the first point of Aries, otherwise known as the vernal, or spring, equinox. The vernal equinox occurs when the sun crosses the equator going from south to north, and an imaginary line drawn from this equatorial crossing through the center of the sun points to the first point of Aries (symbol γ). This is the line of Aries.




Figure 2.4 Prograde and retrograde orbits.




Figure 2.5 The argument of perigee ω and the right ascension of the ascending node Ω .

Six Orbital Elements

- Earth-orbiting artificial satellites are defined by six orbital elements referred to as the *keplerian element set*.
- The semimajor axis a .
- The eccentricity e
 - give the shape of the ellipse.
- A third, the mean anomaly M , gives the position of the satellite in its orbit at a reference time known as the *epoch*.
- A fourth, the argument of perigee ω , gives the rotation of the orbit's perigee point relative to the orbit's line of nodes in the earth's equatorial plane.
- The inclination I
- The right ascension of the ascending node γ
 - Relate the orbital plane's position to the earth.